

VU Research Portal

Complete genome sequence of *Mycobacterium xenopi* type strain RIVM700367

Abdallah, A.M.; Rashid, M.; Adroub, S.A.; Elabdalaoui, H.; Ali, S.; van Soolingen, D.; Bitter, W.; Pain, A.

published in

Journal of Bacteriology
2012

DOI (link to publisher)

[10.1128/JB.00482-12](https://doi.org/10.1128/JB.00482-12)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Abdallah, A. M., Rashid, M., Adroub, S. A., Elabdalaoui, H., Ali, S., van Soolingen, D., Bitter, W., & Pain, A. (2012). Complete genome sequence of *Mycobacterium xenopi* type strain RIVM700367. *Journal of Bacteriology*, 194(12), 3282-3283. <https://doi.org/10.1128/JB.00482-12>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Complete Genome Sequence of *Mycobacterium xenopi* Type Strain RIVM700367

Abdallah M. Abdallah, Mamoon Rashid, Sabir A. Adroub,
Hafida Elabdalaoui, Shahjahan Ali, Dick van Soolingen,
Wilbert Bitter and Arnab Pain
J. Bacteriol. 2012, 194(12):3282. DOI: 10.1128/JB.00482-12.

Updated information and services can be found at:
<http://jb.asm.org/content/194/12/3282>

REFERENCES

These include:

This article cites 13 articles, 6 of which can be accessed free at:
<http://jb.asm.org/content/194/12/3282#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new
articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Complete Genome Sequence of *Mycobacterium xenopi* Type Strain RIVM700367

Abdallah M. Abdallah,^a Mamoon Rashid,^a Sabir A. Adroub,^a Hafida Elabdalaoui,^b Shahjahan Ali,^b Dick van Soolingen,^{c,d} Wilbert Bitter,^e and Arnab Pain^a

Pathogen Genomics Laboratory, Computational Bioscience Research Center, King Abdullah University of Science and Technology, Thuwal-Jeddah, Saudi Arabia^a; Bioscience Core Laboratory, King Abdullah University of Science and Technology, Thuwal-Jeddah, Saudi Arabia^b; Dutch National Institute for Public Health and the Environment, Bilthoven, The Netherlands^c; Department of Clinical Microbiology and Department of Pulmonary Diseases, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands^d; and Department of Medical Microbiology and Infection Control, VU University Medical Center, Amsterdam, The Netherlands^e

***Mycobacterium xenopi* is a slow-growing, thermophilic, water-related *Mycobacterium* species. Like other nontuberculous mycobacteria, *M. xenopi* more commonly infects humans with altered immune function, such as chronic obstructive pulmonary disease patients. It is considered clinically relevant in a significant proportion of the patients from whom it is isolated. We report here the whole genome sequence of *M. xenopi* type strain RIVM700367.**

The genus *Mycobacterium* comprises over 150 known species and 13 subspecies (3), of which nearly one-third have been associated with disease in humans. *Mycobacterium xenopi* is a slow-growing, waterborne, and scotochromogenic mycobacterium. It was first isolated from skin lesions in a clawed frog (*Xenopus laevis*) in 1959 (10). It is a well-known opportunistic pathogen commonly associated with disease in humans, notably, pulmonary infections. Along with *M. avium* and *M. kansasii*, *M. xenopi* represents one of the most common agents of mycobacterial pulmonary infections other than tuberculosis (13). The clinical presentation of the lung infection caused by *M. xenopi* mimics the disease caused by *Mycobacterium tuberculosis* complex (MTC) and *Mycobacterium avium-M. intracellulare* (4). Extrapulmonary and disseminated infections have also been reported, particularly infections involving bone, joints, and muscle (1, 5, 7, 9, 12). However, knowledge on the pathogenesis underlying *M. xenopi* infection is limited. As a first step toward elucidating the molecular basis of *M. xenopi* virulence mechanisms, we have characterized the complete genome sequence of this microorganism. Whole-genome sequencing is also important in order to facilitate more precise identification and rapid diagnosis of *M. xenopi* infections in patients. For instance, on the basis of just limited genomic information, the designation *M. noviomagense* was recently introduced to mark a subgrouping of *M. xenopi* not associated with any clinical importance.

The *M. xenopi* type strain RIVM700367 genome was sequenced by a whole-genome shotgun strategy using the Illumina HiSeq2000 platform. The genome assembly was performed using the *de novo* assembler Velvet (14) and resulted in 117 contigs with an N₅₀ of 154,665 bp, comprising in total 4,434,836 bp. The overall GC content of the chromosome amounted to 66.12%. The genome annotation was performed using the NCBI Prokaryotic Genomes Automatic Annotation Pipeline (PGAAP). The entire genome of *M. xenopi* was predicted to encode 4,282 coding sequences (CDSs), five sets of rRNA operons, and 46 tRNA-encoding genes. It was possible to assign a biological function to 64% (2,749) of the coding sequences on the *M. xenopi* chromosome.

The RAST server annotation pipeline (2) showed that *M. xenopi* has the highest similarity with *M. avium* subsp. *paratuberculosis*

(6), compared to all mycobacteria with complete genome sequences currently available. Furthermore, our characterization of the *M. xenopi* genome showed it to be smaller (4.4 Mb) than the genome of *M. avium* subsp. *paratuberculosis* (4.8 Mb) and to carry fewer genes (4,333 versus 4,399).

Previous studies hypothesized about the evolution of pathogenic nontuberculous mycobacteria from nonpathogenic forms (8, 11). In this regard, the presence of genes involved in virulence and pathogenicity are of particular interest. Careful inspection of the *M. xenopi* genome showed that this genome encodes several genes associated with pathogenicity in other mycobacteria, such as PE and PPE proteins, mammalian cell entry (MCE) family proteins, and esterases/lipases. Interestingly, the important virulence locus *esx-1* is missing in *M. xenopi*, similar to what is observed for species of the *M. avium* complex. Further manual annotation of the genome and functional analysis will provide more information regarding the possible evolution of the pathogenicity of *M. xenopi*.

Nucleotide sequence accession numbers. The results from this whole-genome shotgun project have been deposited with DDBJ/EMBL/GenBank under accession number [AJFI000000000](https://www.ncbi.nlm.nih.gov/nuccore/AJFI000000000). The version described in this paper is the first version, [AJFI000000000](https://www.ncbi.nlm.nih.gov/nuccore/AJFI000000000).

ACKNOWLEDGMENTS

This work was supported by faculty funding to A.P. by the King Abdullah University of Science and Technology. M.R. was supported by a SABIC postdoctoral fellowship.

We thank the Dutch National Institute for Public Health and the Environment (RIVM) for providing the strain. We gratefully acknowledge

Received 26 March 2012 Accepted 9 April 2012

Address correspondence to Abdallah M. Abdallah, Abdallah.Abdallah@kaust.edu.sa, or Arnab Pain, arnab.pain@kaust.edu.sa.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.00482-12

the support of Eveline Weerdenbrug and Roy Ummels at the Department of Medical Microbiology and Infection Control, VU University Medical Centre, Amsterdam, The Netherlands, and Thomas Dan Otto from The Wellcome Trust Sanger Institute, Cambridge, United Kingdom.

REFERENCES

1. Astagneau P, et al. 2001. *Mycobacterium xenopi* spinal infections after disvertebral surgery: investigation and screening of a large outbreak. *Lancet* 358:747–751.
2. Aziz RK, et al. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. doi:10.1186/1471-2164-9-75.
3. Euzéby JP. 2012. List of prokaryotic names with standing in nomenclature, genus *Mycobacterium*. Société de Bactériologie Systématique et Vétérinaire, Blagnac, France. <http://www.bacterio.cict.fr/>.
4. Griffith DE, et al. 2007. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am. J. Resp. Crit. Care* 175:367–416.
5. Huang DB, Gupta A, Gupta G. 2003. Psoas abscess due to *Mycobacterium xenopi* infection. *Am. J. Med.* 114:247–249.
6. Li LL, et al. 2005. The complete genome sequence of *Mycobacterium avium* subspecies *paratuberculosis*. *Proc. Natl. Acad. Sci. U. S. A.* 102:12344–12349.
7. Marks J, Cook J, Pringle JAS. 1975. Bone abscess due to *Mycobacterium xenopi*. *Tubercle* 56:157–159.
8. Ripoll F, et al. 2009. Non mycobacterial virulence genes in the genome of the emerging pathogen *Mycobacterium abscessus*. *Plos One* 4(6):e5660. doi:10.1371/journal.pone.0005660.
9. Salliot C, et al. 2006. Arthritis due to *Mycobacterium xenopi*: a retrospective study of 7 cases in France. *Clin. Infect. Dis.* 43:987–993.
10. Schwabacher H. 1959. A strain of *Mycobacterium* isolated from skin lesions of a cold-blooded animal, *Xenopus laevis*, and its relation to atypical acid-fast bacilli occurring in man. *J. Hyg.* 57:57–67.
11. Stinear TP, et al. 2008. Insights from the complete genome sequence of *Mycobacterium marinum* on the evolution of *Mycobacterium tuberculosis*. *Genome Res.* 18:729–741.
12. van Ingen J, et al. 2008. *Mycobacterium xenopi* clinical relevance and determinants, the Netherlands. *Emerg. Infect. Dis.* 14:385–389.
13. Wolinsky E. 1992. Mycobacterial diseases other than tuberculosis. *Clin. Infect. Dis.* 15:1–12.
14. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.